

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : G01N 33/543, B01L 3/02, G01N 33/569, 33/573, C12Q 1/18		A1	(11) International Publication Number: <b>WO 00/14540</b>
			(43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: PCT/IB98/01399		(81) Designated States: AU, BR, CA, CN, IL, JP, KR, MX, NZ, PL, RU, SG, TR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 8 September 1998 (08.09.98)			
(71) Applicant (for all designated States except US): TIBOTEC N.V. [BE/BE]; Intercity Business Park, Generaal de Wittelaan L11 B3, B-2800 Mechelen (BE).		Published With international search report.	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): PAUWELS, Rudi, Wilfried, Jan [BE/BE]; Berentrodendreef 27, B-2820 Bonheiden (BE). ROELANT, Christiaan, Hubert, Simon [BE/BE]; 's Hertogenlaan 115, B-3000 Leuven (BE). VAN ACKER, Koenraad, Lodewijk, August [BE/BE]; Hollebeek 8, B-9140 Temse (BE).			
(74) Agent: RYAN, Anne; Anne Ryan & Co., 60 Northumberland Road, Ballsbridge, Dublin 4 (IE).			

(54) Title: METHOD FOR THE RAPID SCREENING OF ANALYTES



(57) Abstract

A method for the rapid screening of analytes, such as potential drug candidates, comprises the steps of applying a plurality of analytes to be screened onto one or more solid support(s) (61) such that the analytes remain isolated from one another; contacting said analyte-carrying solid support(s) (61) with targets provided in a semi-solid or liquid medium, whereby said analytes are released from the solid support(s) (61) to the targets; and measuring analyte-target interactions. This method allows for the manipulation of thousands of different analytes simultaneously. When the analyte is applied to the solid support (61) it can diffuse thereon so as to produce a concentration gradient and serial dilution of analyte if a dose response curve for a candidate drug is required. The method described can be readily automated.